

The Cosmic Coincidence as a Temporal Selection Effect Produced by the Age Distribution of Terrestrial Planets in the Universe

Charles H. Lineweaver¹ & Chas A. Egan^{1,2}

¹ *Planetary Science Institute, Research School of Astronomy and Astrophysics & Research School of Earth Sciences, Australian National University, Canberra, ACT, Australia*

² *Department of Astrophysics, School of Physics, University of New South Wales, Sydney, NSW 2052, Australia*

charley@mso.anu.edu.au

ABSTRACT

The energy densities of matter and the vacuum are currently observed to be of the same order of magnitude: ($\Omega_{m_o} \approx 0.3$) \sim ($\Omega_{\Lambda_o} \approx 0.7$). The cosmological window of time during which this occurs is relatively narrow. Thus, we are presented with the cosmological coincidence problem: Why, just now, do these energy densities happen to be of the same order? Here we show that this apparent coincidence can be explained as a temporal selection effect produced by the age distribution of terrestrial planets in the Universe. We find a large ($\sim 68\%$) probability that observations made from terrestrial planets will result in finding Ω_m at least as close to Ω_Λ as we observe today. Hence, we, and any observers in the Universe who have evolved on terrestrial planets, should not be surprised to find $\Omega_{m_o} \sim \Omega_{\Lambda_o}$. This result is relatively robust if the time it takes an observer to evolve on a terrestrial planet is less than ~ 10 Gyr.

Subject headings: Terrestrial Planets, Cosmology, Planets, Cosmic Coincidence

1. Is the Cosmic Coincidence Remarkable or Insignificant?

1.1. Dicke's argument

Dirac (1937) pointed out the near equality of several large fundamental dimensionless numbers of the order 10^{40} . One of these large numbers varied with time since it depended on the age of the Universe. Thus there was a limited time during which this near equality would hold. Under the assumption that observers could exist at any time during the history of the

Universe, this large number coincidence could not be explained in the standard cosmology. This problem motivated Dirac (1938) and Jordan (1955) to construct an ad hoc new cosmology. Alternatively, Dicke (1961) proposed that our observations of the Universe could only be made during a time interval after carbon had been produced in the Universe and before the last stars stop shining. Dicke concluded that this temporal observational selection effect – even one so loosely delimited – could explain Dirac’s large number coincidence without invoking a new cosmology.

Here, we construct a similar argument to address the cosmic coincidence: Why just now do we find ourselves in the relatively brief interval during which $\Omega_m \sim \Omega_\Lambda$. The temporal constraints on observers that we present are more empirical and specific than those used in Dicke’s analysis, but the reasoning is similar. Our conclusion is also similar: a temporal observational selection effect can explain the apparent cosmic coincidence. That is, given the evolution of Ω_Λ and Ω_m in our Universe, most observers in our Universe who have emerged on terrestrial planets will find $\Omega_\Lambda \sim \Omega_m$. Rather than being an unusual coincidence, it is what one should expect.

There are two distinct problems associated with the cosmological constant (Weinberg 2000; Garriga and Vilenkin 2001; Steinhardt 2003). One is the coincidence problem that we address here. The other is the smallness problem and has to do with the observed energy density of the vacuum, ρ_Λ . Why is ρ_Λ so small compared to the $\sim 10^{120}$ times larger value predicted by particle physics? Anthropic solutions to this problem invoke a multiverse and argue that galaxies would not form and there would be no life in a Universe, if ρ_Λ were larger than ~ 100 times its observed value (Weinberg 1987; Martel et al. 1998; Garriga and Vilenkin 2001; Pogosian and Vilenkin 2007). Such explanations for the smallness of ρ_Λ do not explain the temporal coincidence between the time of our observation and the time of the near-equality of Ω_m and Ω_Λ . Here we address this temporal coincidence in our Universe, not the smallness problem in a multiverse.

1.2. Evolution of the Energy Densities

Given the currently observed values for H_o and the energy densities Ω_{r_o} , Ω_{m_o} and Ω_{Λ_o} in the Universe (Spergel et al. 2006; Seljak et al. 2006), the Friedmann equation tells us the evolution of the scale factor a , and the evolution of these energy densities. These are plotted in Fig. 1. The history of the Universe can be divided chronologically into four distinct periods each dominated by a different form of energy: initially the false vacuum energy of inflation dominates, then radiation, then matter, and finally vacuum energy. Currently the Universe is making the transition from matter domination to vacuum energy domination. In an ex-




Fig. 1.— The time dependence of the densities of the major components of the Universe. Given the observed Hubble constant, H_o and energy densities in the Universe today, Ω_{r_o} , Ω_{m_o} , Ω_{Λ_o} (radiation, matter and cosmological constant), we use the Friedmann equation to plot the temporal evolution of the components of the Universe in g/cm³ (top panel), or normalized to the time-dependent critical density $\rho_{\text{crit}} = \frac{3H(t)^2}{8\pi G}$ (bottom panel). We assume an epoch of inflation at $\sim 10^{-35}$ seconds after the big bang and a false vacuum energy density $\rho_{\Lambda \text{ inf}}$ between the Planck scale and t_{GUT} . See Table 1 and Appendix A for details.

panding Universe, with an initial condition $\Omega_m > \Omega_\Lambda > 0$, there will be some epoch in which $\Omega_m \sim \Omega_\Lambda$, since ρ_m is decreasing as $\propto 1/a^3$ while ρ_Λ is a constant (see top panel of Fig. 1 and Appendix A). Figure 1 also shows that the transition from matter domination to vacuum energy domination is occurring now. When we view this transition in the context of the time evolution of the Universe (Fig. 2) we are presented with the cosmic coincidence problem: Why just now do we find ourselves at the relatively brief interval during which this transition happens? Carroll (2001*b,a*) and Dodelson et al. (2000) find this coincidence to be a remarkable result that is crucial to understand. The cosmic coincidence problem is often regarded as an important unsolved problem whose solution may help unravel the nature of dark energy (Turner 2001; Carroll 2001*a*). The coincidence problem is one of the main motivations for the tracker potentials of quintessence models (Caldwell et al. 1998; Steinhardt et al. 1999; Zlatev et al. 1999; Wang et al. 2000; Dodelson et al. 2000; Armendariz-Picon et al. 2000; Guo and Zhang 2005). In these models the cosmological constant is replaced by a more generic form of dark energy in which Ω_m and Ω_Λ are in near-equality for extended periods of time. It is not clear that these models successfully explain the coincidence without fine-tuning (see Weinberg 2000; Bludman 2004).

The interpretation of the observation $\Omega_{m_o} \sim \Omega_{\Lambda_o}$ as a remarkable coincidence in need of explanation depends on some assumptions that we quantify to determine how surprising this apparent coincidence is. We begin this quantification by introducing a time-dependent proximity parameter,

$$r = \min \left[\frac{\Omega_\Lambda}{\Omega_m}, \frac{\Omega_m}{\Omega_\Lambda} \right] \quad (1)$$

which is equal to one when $\Omega_m = \Omega_\Lambda$ and is close to zero when $\Omega_m \gg \Omega_\Lambda$ or $\Omega_m \ll \Omega_\Lambda$. The current value is $r_o \approx 0.4$. In Figure 2 we plot r as a function of $\log(\text{scale factor})$ in the upper panel and as a function of $\log(\text{time})$ in the lower panel. These logarithmic axes allow a large dynamic range that makes our existence at a time when $r \sim 1$, appear to be an unlikely coincidence. This appearance depends on the implicit assumption that we could make cosmological observations at any time with equal likelihood. More specifically, the implicit assumption is that the *a priori* probability distribution P_{obs} , of the times we could have made our observations, is uniform in $\log t$, or $\log a$, over the interval shown.

Our ability to quantify the significance of the coincidence depends on whether we assume that P_{obs} is uniform in time, $\log(\text{time})$, scale factor or $\log(\text{scale factor})$. That is, our result depends on whether we assume: $P_{obs}(t) = \text{constant}$, $P_{obs}(\log t) = \text{constant}$, $P_{obs}(a) = \text{constant}$ or $P_{obs}(\log a) = \text{constant}$. These are the most common possibilities, but there are others. For a discussion of the relative merits of log and linear time scales and implicit uniform priors see Section 3.3 and Jaynes (1968).

In Fig. 3 we plot $r(t)$ on an axis linear in time where the implicit assumption is that the

a priori probability distribution of our existence is uniform in t over the intervals $[0, 100]$ Gyr (top panel) and $[0, 13.8]$ Gyr (bottom panel). The bottom panel shows that the observation $r > 0.4$ could have been made anytime during the past 7.8 Gyr. Thus, our current observation that $r_o \approx 0.4$, does not appear to be a remarkable coincidence. Whether this most recent 7.8 Gyr period is seen as “brief” (in which case there is an unlikely coincidence in need of explanation) or “long” (in which case there is no coincidence to explain) depends on whether we view the issue in log time (Fig. 2) or linear time (Fig. 3).

A large dynamic range is necessary to present the fundamental changes that occurred in the very early Universe, e.g., the transitions at the Planck time, inflation, baryogenesis, nucleosynthesis, recombination and the formation of the first stars. Thus a logarithmic time axis is often preferred by early Universe cosmologists because it seems obvious, from the point of view of fundamental physics, that the cosmological clock ticks logarithmically. This defensible view and the associated logarithmic axis gives the impression that there is a coincidence in need of an explanation. The linear time axis gives a somewhat different impression. Evidently, deciding whether a coincidence is of some significance or only an accident is not easy (Peebles 1999). We conclude that although the importance of the cosmic coincidence problem is subjective, it is important enough to merit the analysis we perform here.

The interpretation of the observation $\Omega_{m_o} \sim \Omega_{\Lambda_o}$ as a coincidence in need of explanation depends on the *a priori* (not necessarily uniform) probability distribution of our existence. That is, it depends on when cosmological observers can exist. We propose that the cosmic coincidence problem can be more constructively evaluated by replacing these uninformed uniform priors with the more realistic assumption that observers capable of measuring cosmological parameters are dependent on the emergence of high density regions of the Universe called terrestrial planets, which require non-trivial amounts of time to form – and that once these planets are in place, the observers themselves require non-trivial amounts of time to evolve.

In this paper we use the age distribution of terrestrial planets estimated by Lineweaver (2001) to constrain when in the history of the Universe, observers on terrestrial planets can exist. In Section 2, we briefly describe this age distribution (Fig. 4) and show how it limits the existence of such observers to an interval in which $\Omega_m \sim \Omega_\Lambda$ (Fig. 5). Using this age distribution as a temporal selection function, we compute the probability of an observer on a terrestrial planet observing $r \geq r_o$ (Fig. 6). In Section 3 we discuss the robustness of our result and find (Fig. 7) that this result is relatively robust if the time it takes an observer to evolve on a terrestrial planet is less than ~ 10 Gyr. In Section 4 we discuss and summarize our results, and compare it to previous work to resolve the cosmic coincidence




Fig. 2.— Plot of the proximity factor r (see Eq. 1). When the matter and vacuum energy densities of the Universe are the same, $\Omega_m = \Omega_\Lambda$, we have $r = 1$. We currently observe $\Omega_{m_0} \sim \Omega_{\Lambda_0}$ and thus, $r \sim 1$. Our existence now when $r \sim 1$ appears to be an unlikely cosmic coincidence when the x axis is logarithmic in the scale factor (top panel) or logarithmic in time (bottom panel). In the top panel, following Carroll (2001*b*), we have chosen a range of scale factors with “Now” midway between the scale factor at the Planck time and the scale factor at the inverse Planck time [$a_{Planck} < a < a_{Planck}^{-1}$]. The brief epoch shown in grey between the thin vertical lines is the epoch during which $r > r_o$ (where $r_o \approx 0.4$ is the currently observed value). In the bottom panel the range shown on the x axis is [$t_{Planck} < t < 10^{22}$] seconds. The Planck time and Planck scale provide reasonably objective lower time limits. The upper limits are somewhat arbitrary but contribute to the impression that $r \approx 0.4 \sim 1$ is an unlikely coincidence.




Fig. 3.— Plot of the proximity factor r , as in the previous figure, but plotted here with a linear rather than a logarithmic time axis. The condition $r > r_o \approx 0.4$ does not seem as unlikely as in the previous figure. The range of time plotted also affects this appearance; with the $[0, 100]$ Gyr range of the top panel, the time interval highlighted in grey where $r > r_o$, appears narrow and relatively unlikely. In contrast, the $[0, 13.8]$ Gyr range of the bottom panel seems to remove the appearance of $r > r_o$ being an unlikely coincidence in need of explanation; for the first ~ 6 Gyrs we have $r < r_o$ while in the subsequent 7.8 Gyr we have $r > r_o$. How can $r > r_o$ be an unlikely coincidence when it has been true for most of the history of the Universe?

problem (Garriga et al. 2000; Bludman and Roos 2001).

2. How We Compute the Probability of Observing $\Omega_m \sim \Omega_\Lambda$

2.1. The Age Distribution of Terrestrial Planets and New Observers

The mass histogram of detected extrasolar planets peaks at low masses: $dN/dM \propto M^{-1.7}$, suggesting that low mass planets are abundant (Lineweaver and Grether 2003). Terrestrial planet formation may be a common feature of star formation (Wetherill 1996; Chyba 1999; Ida and Lin 2005). Whether terrestrial planets are common or rare, they will have an age distribution proportional to the star formation rate – modified by the fact that in the first ~ 2 billion years of star formation, metallicities are so low that the material for terrestrial planet formation will not be readily available. Using these considerations, Lineweaver (2001) estimated the age distribution of terrestrial planets – how many Earths are produced by the Universe per year, per Mpc^3 (Figure 4). If life emerges rapidly on terrestrial planets (Lineweaver and Davis 2002) then this age distribution is the age distribution of biogenesis in the Universe. However, we are not just interested in any life; we would like to know the distribution in time of when independent observers first emerge and are able to measure Ω_m and Ω_Λ , as we are able to do now. If life originates and evolves preferentially on terrestrial planets, then the Lineweaver (2001) estimate of the age distribution of terrestrial planets is an *a priori* input which can guide our expectations of when we (as members of a hypothetical group of terrestrial-planet-bound observers) could have been present in the Universe. It takes time (if it happens at all) for life to emerge on a new terrestrial planet and evolve into cosmologists who can observe Ω_m and Ω_Λ . Therefore, to obtain the age distribution of new independent observers able to measure the composition of the Universe for the first time, we need to shift the age distribution of terrestrial planets by some characteristic time, Δt_{obs} required for observers to evolve. On Earth, it took $\Delta t_{obs} \sim 4$ Gyr for this to happen. Whether this is characteristic of life elsewhere in the Universe is uncertain (Carter 1983; Lineweaver and Davis 2003). For our initial analysis we use $\Delta t_{obs} = 4$ Gyr as a nominal time to evolve observers. In Section 3.1 we allow Δt_{obs} to vary from 0-12 Gyr to see how sensitive our result is to these variations. Fig. 4 shows the age distribution of terrestrial planet formation in the Universe shifted by $\Delta t_{obs} = 4$ Gyr. This curve, labeled “ P_{obs} ” is a crude prior for the temporal selection effect of when independent observers can first measure r . Thus, if the evolution of biological equipment capable of doing cosmology takes about $\Delta t_{obs} \sim 4$ Gyr, the “ P_{obs} ” in Fig. 4 shows the age distribution of the first cosmologists on terrestrial planets able to look at the Universe and determine the overall energy budget, just as we have recently been able to do.




Fig. 4.— The terrestrial planet formation rate $PFR(t)$, derived in Lineweaver (2001) is an estimate of the age distribution of terrestrial planets in the Universe and is shown here as a thin solid line. Estimated uncertainty is given by the thin dashed lines. To allow time for the evolution of observers on terrestrial planets, we shift this distribution by Δt_{obs} to obtain an estimate of the age distribution of observers: $P_{obs}(t) = PFR(t - \Delta t_{obs})$ (thick solid line). The grey band represents the error estimate on $P_{obs}(t)$ which is the shifted error estimates on $PFR(t)$. In the case shown here $\Delta t_{obs} = 4$ Gyr, which is how long it took life on Earth to emerge, evolve and be able to measure the composition of the Universe. To obtain the numerical values on the y axis, we have followed Lineweaver (2001) and assumed that one out of one hundred stars is orbited by a terrestrial planet. We have smoothly extrapolated the $PFR(t)$ of Lineweaver (2001) into the future. This time dependence and our subsequent analysis does not depend on whether the probability for terrestrial planets to produce observers is high or low.




Fig. 5.— Zoom-in of the portion of Fig. 1 between 1 and 100 billion years after the big bang, containing the relatively narrow window of time in which $\Omega_m \sim \Omega_\Lambda$. The 99 Gyr time interval displayed here is indicated in Fig. 1 by the small grey rectangle above the “Now” label. The proximity parameter $r(t)$ (Eq. 1, Figs. 2 & 3) is superimposed. The thin solid line shows the age distribution of terrestrial planets in the Universe while the thick solid line is the lateral displacement of this distribution by $\Delta t_{obs} = 4$ Gyr. These distributions were presented in Fig. 4, but here the time axis is logarithmic. We interpret P_{obs} as the frequency distribution of new observers able to measure Ω_m and Ω_Λ for the first time. Since $r(t)$ peaks at about the same time as $P_{obs}(t)$, large values of r will be observed more often than small values.

2.2. The Probability of Observing $\Omega_m \sim \Omega_\Lambda$.

In Fig. 5 we zoom into the portion of Fig. 1 containing the relatively narrow window of time in which $\Omega_m \sim \Omega_\Lambda$. We plot $r(t)$ to show where $r \sim 1$ and we also plot the age distribution of planets and the age distribution of recently emerged cosmologists from Fig. 4. The white area under the thick $P_{obs}(t)$ curve provides an estimate of the time distribution of new observers in the Universe. We interpret $P_{obs}(t)$ as the probability distribution of the times at which new, independent observers are able to measure r for the first time.

Lineweaver (2001) estimated that the Earth is relatively young compared to other terrestrial planets in the Universe. It follows under the simple assumptions of our analysis that most terrestrial-planet-bound observers will emerge earlier than we have. We compute the fraction f of observers who have emerged earlier than we have,

$$f = \frac{\int_0^{t_o} P_{obs}(t) dt}{\int_0^\infty P_{obs}(t) dt} \approx 68\% \quad (2)$$

and find that 68% emerge earlier while 32% emerge later. These numbers are indicated in Fig. 5.

2.3. Converting $P_{obs}(t)$ to $P_{obs}(r)$

We have an estimate of the distribution in time of observers, $P_{obs}(t)$, and we have the proximity parameter $r(t)$. We can then convert these to a probability $P_{obs}(r)$, of observed values of r . That is, we change variables and convert the t -dependent probability to an r -dependent probability: $P_{obs}(t) \rightarrow P_{obs}(r)$. We want the probability distribution of the r values first observed by new observers in the Universe. Let the probability of observing r in the interval dr be $P_{obs}(r)dr$. This is equal to the probability of observing t in the interval dt , which is $P_{obs}(t)dt$

Thus,

$$P_{obs}(r) dr = P_{obs}(t) dt \quad (3)$$

or equivalently

$$P_{obs}(r) = \frac{P_{obs}(t)}{dr/dt} \quad (4)$$

where $P_{obs}(t) = PFR(t - \Delta t_{obs})$ is the temporally shifted age distribution of terrestrial planets and dr/dt is the slope of $r(t)$. Both are shown in Fig. 5. The distribution $P_{obs}(r)$ is shown in Fig. 6 along with the upper and lower confidence limits on $P_{obs}(r)$ obtained by

inserting the upper and lower confidence limits of $P_{obs}(t)$ (denoted “ P^+ ” and “ P^- ” in Fig. 4), into Eq. 4 in place of $P_{obs}(t)$.

The probability of observing $r > r_o$ is,

$$P(r > r_o) = \int_{r_o}^1 P_{obs}(r) dr = \int_{t'}^{t_o} P_{obs}(t) dt \approx 68\% \quad (5)$$

where t' is the time in the past when r was equal to its present value, i.e., $r(t') = r(t_o) = r_o \approx 0.4$. We have $t' = 6$ Gyr and $t_o = 13.8$ Gyr (see bottom panel of Fig. 3). This integral is shown graphically in Fig. 6 as the hatched area underneath the “ $P_{obs}(r)$ ” curve, between $r = r_o$ and $r = 1$. We interpret this as follows: of all observers that have emerged on terrestrial planets, 68% will emerge when $r > r_o$ and thus will find $r > r_o$. The 68% from Eq. 2 is only the same as the 68% from Eq. 5 because all observers who emerge earlier than we did, did so more recently than 7.8 billion years ago and thus, observe $r > r_o$ (Fig. 5).




Fig. 6.— Probability of new observers on terrestrial planets observing a given r (Eq. 4). Given our estimate of the age distribution of new cosmologists in the Universe $P_{obs}(t)$, the probability of observing Ω_m and Ω_Λ as close together as they are, or closer, is the integral given in Eq. (5), shown here as the hashed area labeled 68%. The dashed lines labeled P^+ and P^- are from replacing $P_{obs}(t)$ in Eq. 4 with the curves labeled P^+ and P^- in Fig. 4.

We obtain estimates of the uncertainty on this 68% estimate by computing analogous

integrals underneath the curves labeled P^+ and P^- in Fig. 6. These yield 82% and 59% respectively. Thus, under the assumptions made here, $68_{-10}^{+14}\%$ of the observers in the Universe will find Ω_Λ and Ω_m even closer to each other than we do. This suggests that a temporal selection effect due to the constraints on the emergence of observers on terrestrial planets provides a plausible solution to the cosmic coincidence problem. If observers in our Universe evolve predominantly on Earth-like planets (see the “principle of mediocrity” in Vilenkin (1995*b*)), we should not be surprised to find ourselves on an Earth-like planet and we should not be surprised to find $\Omega_{\Lambda_o} \sim \Omega_{m_o}$.

3. How Robust is this 68% Result?

3.1. Dependence on the timescale for the evolution of observers

A necessary delay, required for the biological evolution of observing equipment – e.g. brains, eyes, telescopes, makes the observation of recent biogenesis unobservable (Lineweaver and Davis 2002, 2003). That is, no observer in the Universe can wake up to observerhood and find that their planet is only a few hours old. Thus, the timescale for the evolution of observers, $\Delta t_{obs} > 0$.

Our $68_{-10}^{+14}\%$ result was calculated under the assumption that evolution from a new terrestrial planet to an observer takes $\Delta t_{obs} \sim 4$ Gyr. To determine how robust our result is to variations in Δt_{obs} , we perform the analysis of Sec. 2 for $0 < \Delta t_{obs} < 12$ Gyr. The results are shown in Fig. 7. Our $68_{-10}^{+14}\%$ result is the data point plotted at $\Delta t_{obs} = 4$ Gyr. If life takes ~ 0 Gyr to evolve to observerhood, once a terrestrial planet is in place, $P_{obs}(t) \approx PFR(t)$ and 55% of new cosmologists would observe an r value larger than the $r_o \approx 0.4$ that we actually observe today. If observers typically take twice as long as we did to evolve ($\Delta t_{obs} \sim 8$ Gyr), there is still a large chance ($\sim 30\%$) of observing $r > r_o$. If $\Delta t_{obs} > 11$ Gyr, $P_{obs}(t)$ in Fig. 5 peaks substantially after $r(t)$ peaks, and the percentage of cosmologists who see $r > r_o$, is close to zero (Eq. 5). Thus, if the characteristic time it takes for life to emerge and evolve into cosmologists is $\Delta t_{obs} \lesssim 10$ Gyr, our analysis provides a plausible solution to the cosmic coincidence problem.

The Sun is more massive than 94% of all stars. Therefore 94% of stars live longer than the $t_\odot \approx 10$ Gyr main sequence lifetime of the Sun. This is mildly anomalous and it is plausible that the Sun’s mass has been anthropically selected. For example, perhaps stars as massive as the Sun are needed to provide the UV photons to jump start and energize the molecular evolution that leads to life. If so, then ~ 10 Gyr is a rough upper limit to the amount of time a terrestrial planet with simple life has to produce observers. Even if the

characteristic time for life to evolve into observers is much longer than 10 Gyr, as concluded by Carter (1983), this UV requirement that life-hosting stars have main sequence lifetimes $\lesssim 10$ Gyr would lead to the extinction of most extraterrestrial life before it can evolve into observers. This would lead to observers waking to observerhood to find the age of their planet to be a large fraction of the main sequence lifetime of their star; the time they took to evolve would satisfy $\Delta t_{obs} \lesssim 10$ Gyr, and they would observe that $r \sim 1$ and that other observers are very rare. Such is our situation.

If we assume that we are typical observers (Vilenkin 1995*a,b*, 1996*a,b*) and that the coincidence problem must be resolved by an observer selection effect (Bostrom 2002), then we can conclude that the typical time it takes observers to evolve on terrestrial planets is less than 10 Gyr ($\Delta t_{obs} < 10$ Gyr).

3.2. Dependence on the age distribution of terrestrial planets

The $P_{obs}(t)$ used here (Fig. 5) is based on the star formation rate (SFR) computed in Lineweaver (2001). There is general agreement that the SFR has been declining since redshifts $z \sim 2$. Current debate centers around whether that decline has only been since $z \sim 2$ or whether the SFR has been declining from a much higher redshift (Lanzetta et al. 2002; Hopkins and Beacom 2006; Nagamine et al. 2006; Thompson et al. 2006). Since Lineweaver (2001) assumed a relatively high value for the SFR at redshifts above 2, this led to a relatively high estimate of the metallicity of the Universe at $z \sim 2$, which corresponds to a relatively short delay (~ 2 Gyr) between the big bang and the first terrestrial planets. For the purposes of this analysis, the early-SFR-dependent uncertainty in the ~ 2 Gyr delay is degenerate with, but much smaller than, the uncertainty of Δt_{obs} . Thus the variations of Δt_{obs} discussed above subsume the SFR-dependent uncertainty in $P_{obs}(t)$.

3.3. Dependence on Measure

In Figs. 2 & 3 we illustrated how the importance of the cosmic coincidence depends on the range over which one assumes that the observation of r could have occurred. This involved choosing the range Δx shown on the x axis in Figs. 2 & 3. We also showed how the apparent significance of the coincidence depended on how one expressed that range, i.e., logarithmic in Fig. 2 and linear in Fig. 3. The coincidence seems most compelling when Δx is the largest and the problem is presented on a logarithmic x axis. This dependence is a specific example of a “measure” problem (Aguirre and Tegmark 2005; Aguirre et al. 2006).




Fig. 7.— Percentage of cosmologists who see $r > r_o$ as a function of the time Δt_{obs} , it takes observers to evolve on a terrestrial planet. Since we have only vague notions about how long it takes observers to evolve on a planet, we vary Δt_{obs} between 0 and 12 billion years and show how the probability $P(r > r_o)$ of observing $r > r_o$ (Eq. 5) varies as a function of Δt_{obs} . The $68^{+14}_{-10}\%$ point plotted is the result from Fig. 6 where $\Delta t_{obs} = 4$ Gyr. If $\Delta t_{obs} = 0$, we use the thin solid line in Fig. 5 as $P_{obs}(t)$ rather than the thick solid line and we obtain 55%.

The measure problem is illustrated in Fig. 8, where we plot four different uniform distributions of observers on a linear time axis. In Panel *a*) $P_{obs}(t) = \text{constant}$. That is, we assume that observers could find themselves anywhere between $t_{rec} = 380,000$ yr and 100 Gyr after the big bang, with uniform probability (dark grey). In *b*), we make the different assumption that observers are distributed uniformly in $\log(t)$ over the same range in time. This means for example that the probability of finding yourself between 0.1 and 1 Gyr is the same as between 1 and 10 Gyr. We plot this as a function of linear time and find that the distribution of observers (dark grey) is highest towards earlier times.

To quantify and explore these dependencies further, in Table 2, we take the duration when $r > r_o$ (call this interval Δx_r) and divide it by various larger ranges Δx (a range of




Fig. 8.— The expected observed value of r depends strongly on the assumed distribution of observers over time t . This figure demonstrates a variety of uniform observer distributions P_{obs} which, if used, result in the cosmic coincidence problem that the observed value of r is unexpectedly high. The P_{obs} that are functions of $\log(a)$ or $\log(t)$ have been normalized to the interval t_{rec} to 100 Gyr. Panel *a*) is the same as the top panel of Fig. 3. The probabilities that an observer would fall within the vertical light grey band ($r > r_o$) in Panels *a, b, c* and *d* are 8%, 7%, 0.2% and 6% respectively, and are given in the first row of Table 2.

time or scale factor). Thus, when the probability $P(r > r_o) = \frac{\Delta x_r}{\Delta x}$ is $\ll 1$, there is a low probability that one would find oneself in the interval Δx_r and the cosmic coincidence is compelling. However, when $P(r > r_o) \sim 1$ the coincidence is not significant.

In the four panels *a, b, c* and *d* of Fig. 8 the probability of us observing $r \geq r_o$ (finding ourselves in the light grey area) is respectively 8%, 7%, 0.2% and 6%. These values are given in the first row of Table 2 along with analogous values when 11 other ranges for Δx are considered. Probabilities corresponding to the four panels of Figs. 2 & 3 are shown in bold in Table 2. Our conclusion is that this simple ratio method of measuring the significance of a coincidence yields results that can vary by many orders of magnitude depending on the range (Δx) and measure (e.g. linear or logarithmic) chosen. The use of the non-uniform $P_{obs}(t)$ shown in Fig. 4 is not subject to these ambiguities in the choice of range and measure.

4. Discussion & Summary

Anthropic arguments to resolve the coincidence problem include Garriga et al. (2000) and Bludman and Roos (2001). Both use a semi-analytical formalism (Gunn and Gott 1972; Press and Schechter 1974; Martel et al. 1998) to compute the number density of objects that collapse into large galaxies. This is then used as a measure of the number density of intelligent observers. Our work complements these semi-analytic models by using observations of the star formation rate to constrain the possible times of observation. Our work also extends this previous work by including the effect of Δt_{obs} , the time it takes observers to evolve on terrestrial planets. This inclusion puts an important limit on the validity of anthropic solutions to the coincidence problem.

Garriga et al. (2000) is probably the work most similar to ours. They take ρ_Λ as a random variable in a multiverse model with a prior probability distribution. For a wide range of ρ_Λ (prescribed by a prior based on inflation theory) they find approximate equality between the time of galaxy formation t_G , the time when Λ starts to dominate the energy density of the Universe t_Λ and now t_o . That is, they find that, within one order of magnitude, $t_G \sim t_\Lambda \sim t_o$. Their analysis is more generic but approximate in that it addresses the coincidence for a variety of values of ρ_Λ to an order of magnitude precision. Our analysis is more specific and empirical in that we condition on our Universe and use the Lineweaver (2001) star-formation-rate-based estimate of the age distribution of terrestrial planets to reach our main result (68%).

To compare our result to that of Garriga et al. (2000), we limit their analysis to the ρ_Λ observed in our Universe ($\rho_\Lambda = 6.7 \times 10^{-30} g/cm^3$) and differentiate their cumulative number

of galaxies which have assembled up to a given time (their Eq. 9). We find a broad time-dependent distribution for galaxy formation which is the analog of our more empirical and narrower (by a factor of 2 or 3) $P_{obs}(t)$.

We have made the most specific anthropic explanation of the cosmic coincidence using the age distribution of terrestrial planets in our Universe and found this explanation fairly robust to the largely uncertain time it takes observers to evolve. Our main result is an understanding of the cosmic coincidence as a temporal selection effect if observers emerge preferentially on terrestrial planets in a characteristic time $\Delta t_{obs} < 10$ Gyr. Under these plausible conditions, we, and any observers in the Universe who have evolved on terrestrial planets, should not be surprised to find $\Omega_{m_o} \sim \Omega_{\Lambda_o}$.

Acknowledgements We would like to thank Paul Francis and Charles Jenkins for helpful discussions. CE acknowledges a UNSW School of Physics post graduate fellowship.

Appendix A: Evolution of Densities

Recent cosmological observations have led to the new standard Λ CDM model in which the density parameters of radiation, matter and vacuum energy are currently observed to be $\Omega_{r_o} \approx 4.9 \pm 0.5 \times 10^{-5}$, $\Omega_{m_o} \approx 0.26 \pm 0.03$ and $\Omega_{\Lambda_o} \approx 0.74 \pm 0.03$ respectively and Hubble's constant is $H_o = 71 \pm 3 \text{ km.s}^{-1} \text{ Mpc}^{-1}$ (Spergel et al. 2006; Seljak et al. 2006).

The energy densities in relativistic particles (“radiation” i.e., photons, neutrinos, hot dark matter), non-relativistic particles (“matter” i.e., baryons, cold dark matter) and in vacuum energy scale differently (Peacock 1999),

$$\rho_i \propto a^{-3(w_i+1)}. \quad (6)$$

Where the different equations of state are, $\rho_i = w_i p$ where $w_{radiation} = 1/3$, $w_{matter} = 0$ and $w_{\Lambda} = -1$ (Linder 1997). That is, as the Universe expands, these different forms of energy density dilute at different rates.

$$\rho_r \propto a^{-4} \quad (7)$$

$$\rho_m \propto a^{-3} \quad (8)$$

$$\rho_{\Lambda} \propto a^0 \quad (9)$$

Given the currently observed values for Ω_r , Ω_m and Ω_{Λ} , the Friedmann equation for a standard flat cosmology tells us the evolution of the scale factor of the Universe, and the history of the energy densities:

$$\left(\frac{\dot{a}}{a}\right)^2 = \frac{8\pi G}{3}(\rho_r + \rho_m + \rho_{\Lambda}) \quad (10)$$

$$= \frac{8\pi G}{3}(\rho_{r_o}a^{-4} + \rho_{m_o}a^{-3} + \rho_{\Lambda}a^0) \quad (11)$$

$$= (\Omega_{r_o}a^{-4} + \Omega_{m_o}a^{-3} + \Omega_{\Lambda_o}a^0) \quad (12)$$

where we have $\rho_{crit} = \frac{3H(t)^2}{8\pi G}$ and $\Omega_i = \frac{\rho_i}{\rho_{crit}}$. The upper panel of Fig. 1 illustrates these different dependencies on scale factor and time in terms of densities while the lower panel shows the corresponding normalized density parameters. A false vacuum energy $\rho_{\Lambda_{inf}}$ is assumed between the Planck scale and the GUT scale. In constructing this density plot and setting a value for $\Omega_{\Lambda_{inf}}$ we have used the constraint that at the GUT scale, all the energy densities add up to $\rho_{\Lambda_{inf}}$ which remains constant at earlier times.

Appendix B: Tables

Table 1: Important Times in the History of the Universe (some used in Table 2)

Event (ref)	Symbol	Time after Big Bang	
		seconds	Gyr
Planck time, beginning of time	t_{Planck}	5.4×10^{-44}	1.7×10^{-60}
end of inflation, reheating, origin of matter, thermalization	t_{reheat}	$[10^{-43}, 10^{-33}]$	$[10^{-60}, 10^{-50}]$
energy scale of Grand Unification Theories (GUT)	t_{GUT}	10^{-33}	10^{-50}
matter-anti-matter annihilation, baryogenesis	$t_{baryogenesis}$	$[10^{-33}, 10^{-12}]$	$[10^{-50}, 10^{-29}]$
electromagnetic and weak nuclear forces diverge	$t_{electroweak}$	10^{-12}	10^{-29}
light atomic nuclei produced	t_{BBN}	$[100, 300]$	$[3, 9] \times 10^{-15}$
radiation-matter equality ¹	t_{r-m}	8.9×10^{11}	2.8×10^{-5}
recombination ¹ (first chemistry)	t_{rec}	1.2×10^{13}	0.38×10^{-3}
first thermal disequilibrium	$t_{1sttherm-dis}$	1.2×10^{13}	0.38×10^{-3}
first stars, Pop III, reionization ¹	$t_{1ststars}$	1×10^{16}	0.4
first terrestrial planets ²	$t_{1stEarths}$	8×10^{16}	2.5
last time r had same value as today	t_{rnow}	1.9×10^{17}	6.1
formation of the Sun, Earth ³	t_{Sun}, t_{Earth}	2.9×10^{17}	9.1
matter- Λ equality ¹	$t_{m-\Lambda}$	3.0×10^{17}	9.4
now	t_o	4.4×10^{17}	13.8
last stars die ⁴	$t_{laststars}$	10^{22}	10^6
protons decay ⁴	$t_{protondecay}$	10^{45}	10^{29}
super massive black holes consume matter ⁴	$t_{blackholes}$	10^{107}	10^{91}
maximum entropy (no gradients to drive life) ⁴	$t_{heatdeath}$	10^{207}	10^{191}

References:

- (1) Spergel et al. 2006, <http://map.gsfc.nasa.gov/>
- (2) Lineweaver 2001
- (3) Allègre et al. 1995
- (4) Adams and Laughlin 1997

Table 2. The probability $P(r > r_o)$ of observing $r > r_o$ assuming a uniform distribution of observers P_{obs} in linear time, $\log(\text{time})$, scale factor and $\log(\text{scale factor})$ within the range Δx listed.

Range Δx ^a		$P(r > r_o)$ [%]			
x_{min}	x_{max}	t	$\log(t)$	a	$\log(a)$
t_{rec}	100 Gyr ^b	8 ^c	7	0.2	6
t_{Planck}	$t_{laststars}$	8×10^{-4}	0.6 ^d	10^{-10^4}	10^{-3}
t_{Planck}	t_o	60 ^c	0.6	50	1
t_{Planck}	t_{Planck}^{-1}	30	0.3	30	0.5 ^d
t_{Planck}	$t_{heatdeath}$	10^{-188}	0.1	$10^{-10^{189}}$	10^{-188}
t_{rec}	$t_{protondecay}$	10^{-26}	1	$10^{-10^{27}}$	10^{-26}
t_{rec}	$t_{blackholes}$	10^{-88}	0.4	$10^{-10^{89}}$	10^{-88}
t_{rec}	$t_{heatdeath}$	10^{-188}	0.2	$10^{-10^{189}}$	10^{-188}
$t_{1ststars}$	$t_{laststars}$	8×10^{-4}	6	10^{-10^4}	10^{-3}
$t_{1ststars}$	$t_{protondecay}$	10^{-26}	1	$10^{-10^{27}}$	10^{-26}
$t_{1ststars}$	$t_{blackholes}$	10^{-88}	0.4	$10^{-10^{89}}$	10^{-88}
$t_{1ststars}$	$t_{heatdeath}$	10^{-188}	0.2	$10^{-10^{189}}$	10^{-188}

^a See Table 1 for the times corresponding to columns 1 and 2.

^b The four values in the top row correspond to Fig. 8.

^c The two values shown in bold in the t column correspond to the two panels of Fig. 3.

^d These values correspond to the two panels of Fig. 2.

REFERENCES

- Adams, F. C. and Laughlin, G. (1997), ‘A dying universe: the long-term fate and evolution of astrophysical objects’, *Reviews of Modern Physics* **69**, 337–372.
- Aguirre, A., Gratton, S. and Johnson, M. C. (2006), ‘Hurdles for Recent Measures in Eternal Inflation’, *ArXiv hep-th/0611221* .
- Aguirre, A. and Tegmark, M. (2005), ‘Multiple universes, cosmic coincidences, and other dark matters’, *Journal of Cosmology and Astro-Particle Physics* **1**, 3–19.
- Allègre, C. J., Manhès, G. and Göpel, C. (1995), ‘The age of the Earth’, *Geochim. Cosmochim. Acta* **59**, 1445–1456.
- Armendariz-Picon, C., Mukhanov, V. and Steinhardt, P. J. (2000), ‘Dynamical Solution to the Problem of a Small Cosmological Constant and Late-Time Cosmic Acceleration’, *Physical Review Letters* **85**, 4438–4441.
- Bludman, S. (2004), ‘Tracking quintessence would require two cosmic coincidences’, *Phys. Rev. D* **69**(12), 122002.
- Bludman, S. A. and Roos, M. (2001), ‘Smooth Energy: Cosmological Constant or Quintessence?’, *ApJ* **547**, 77–81.
- Bostrom, N. (2002), *Anthropic Bias: Observation Selection Effects in Science and Philosophy*, Routledge, London.
- Caldwell, R. R., Dave, R. and Steinhardt, P. J. (1998), ‘Cosmological Imprint of an Energy Component with General Equation of State’, *Physical Review Letters* **80**, 1582–1585.
- Carroll, S. M. (2001a), ‘Dark Energy and the Preposterous Universe’, *SNAP (SuperNova Acceleration Probe) Yellow Book astro-ph/0107571* .
- Carroll, S. M. (2001b), ‘The Cosmological Constant’, *Living Reviews in Relativity* **4**, 1.
- Carter, B. (1983), ‘The anthropic principle and its implications for biological evolution’, *Philosophical Transactions of the Royal Society A* **310**, 347–363.
- Chyba, C. F. (1999), ‘Free Energy for Life on Europa’, *AAS/Division for Planetary Sciences Meeting Abstracts* **31**, 66.09.
- Dicke, R. H. (1961), ‘Dirac’s Cosmology and Mach’s Principle’, *Nature* **192**, 440.
- Dirac, P. A. M. (1937), ‘The cosmological constants’, *Nature* **139**, 323.

- Dirac, P. A. M. (1938), ‘A New Basis for Cosmology’, *Royal Society of London Proceedings Series A* **165**, 199–208.
- Dodelson, S., Kaplinghat, M. and Stewart, E. (2000), ‘Solving the Coincidence Problem: Tracking Oscillating Energy’, *Physical Review Letters* **85**, 5276–5279.
- Garriga, J., Livio, M. and Vilenkin, A. (2000), ‘Cosmological constant and the time of its dominance’, *Phys. Rev. D* **61**(2), 023503.
- Garriga, J. and Vilenkin, A. (2001), ‘Solutions to the cosmological constant problems’, *Phys. Rev. D* **64**(2), 023517.
- Gunn, J. E. and Gott, J. R. I. (1972), ‘On the Infall of Matter Into Clusters of Galaxies and Some Effects on Their Evolution’, *ApJ* **176**, 1.
- Guo, Z.-K. and Zhang, Y.-Z. (2005), ‘Interacting phantom energy’, *Phys. Rev. D* **71**(2), 023501.
- Hopkins, A. M. and Beacom, J. F. (2006), ‘On the Normalization of the Cosmic Star Formation History’, *ApJ* **651**, 142–154.
- Ida, S. and Lin, D. N. C. (2005), ‘Toward a Deterministic Model of Planetary Formation. III. Mass Distribution of Short-Period Planets around Stars of Various Masses’, *ApJ* **626**, 1045–1060.
- Jaynes, E. T. (1968), ‘Prior Probabilities (p16)’, *IEEE Transactions on System Science and Cybernetics* **SSC-4**, 227–241.
- Jordan, P. (1955), *Schwerkraft und Weltall*, Braunschweig, Braunschweig, Germany.
- Lanzetta, K. M., Yahata, N., Pascarella, S., Chen, H.-W. and Fernández-Soto, A. (2002), ‘The Star Formation Rate Intensity Distribution Function: Implications for the Cosmic Star Formation Rate History of the Universe’, *ApJ* **570**, 492–501.
- Linder, E. (1997), *First Principles of Cosmology*, First Principles of Cosmology, by E.V. Linder. Addison-Wesley, Harlow, England, 1997.
- Lineweaver, C. H. (2001), ‘An Estimate of the Age Distribution of Terrestrial Planets in the Universe: Quantifying Metallicity as a Selection Effect’, *Icarus* **151**, 307–313.
- Lineweaver, C. H. and Davis, T. M. (2002), ‘Does the Rapid Appearance of Life on Earth Suggest that Life Is Common in the Universe?’, *Astrobiology* **2**, 293–304.

- Lineweaver, C. H. and Davis, T. M. (2003), ‘On the Nonobservability of Recent Biogenesis’, *Astrobiology* **3**, 241–243.
- Lineweaver, C. H. and Grether, D. (2003), ‘What Fraction of Sun-like Stars Have Planets?’, *ApJ* **598**, 1350–1360.
- Martel, H., Shapiro, P. R. and Weinberg, S. (1998), ‘Likely Values of the Cosmological Constant’, *ApJ* **492**, 29.
- Nagamine, K., Ostriker, J. P., Fukugita, M. and Cen, R. (2006), ‘The History of Cosmological Star Formation: Three Independent Approaches and a Critical Test Using the Extragalactic Background Light’, *ApJ* **653**, 881–893.
- Peacock, J. A. (1999), *Cosmological Physics*, Cosmological Physics, by John A. Peacock, pp. 704. ISBN 052141072X. Cambridge, UK: Cambridge University Press, January 1999.
- Peebles, P. J. E. (1999), ‘Cosmology: Evolution of the cosmological constant’, *Nature* **398**, 25.
- Pogosian, L. and Vilenkin, A. (2007), ‘Anthropic predictions for vacuum energy and neutrino masses in the light of WMAP-3’, *Journal of Cosmology and Astro-Particle Physics* **1**, 25–+.
- Press, W. H. and Schechter, P. (1974), ‘Formation of Galaxies and Clusters of Galaxies by Self-Similar Gravitational Condensation’, *ApJ* **187**, 425–438.
- Seljak, U., Slosar, A. and McDonald, P. (2006), ‘Cosmological parameters from combining the Lyman-alpha forest with CMB, galaxy clustering and SN constraints’, *ArXiv Astrophysics e-prints* .
- Spiegel, D. N., Bean, R., Dore, O., Nolta, M. R., Bennett, C. L., Hinshaw, G., Jarosik, N., Komatsu, E., Page, L., Peiris, H. V., Verde, L., Barnes, C., Halpern, M., Hill, R. S., Kogut, A., Limon, M., Meyer, S. S., Odegard, N., Tucker, G. S., Weiland, J. L., Wollack, E. and Wright, E. L. (2006), ‘Wilkinson Microwave Anisotropy Probe (WMAP) Three Year Results: Implications for Cosmology’, *astro-ph/0603449* <http://lambda.gsfc.nasa.gov/product/map/dr2/params/lcdm-all.cfm> .
- Steinhardt, P. J. (2003), ‘A quintessential introduction to dark energy’, *Royal Society of London Philosophical Transactions Series A* **361**, 2497–2513.
- Steinhardt, P. J., Wang, L. and Zlatev, I. (1999), ‘Cosmological tracking solutions’, *Phys. Rev. D* **59**(12), 123504.

- Thompson, R. I., Eisenstein, D., Fan, X., Dickinson, M., Illingworth, G. and Kennicutt, Jr., R. C. (2006), ‘Star Formation History of the Hubble Ultra Deep Field: Comparison with the Hubble Deep Field-North’, *ApJ* **647**, 787–798.
- Turner, M. S. (2001), ‘Dark Energy’, *Nuclear Physics B Proceedings Supplements* **91**, 405–409.
- Vilenkin, A. (1995*a*), ‘Making predictions in an eternally inflating universe’, *Phys. Rev. D* **52**, 3365–3374.
- Vilenkin, A. (1995*b*), ‘Predictions from Quantum Cosmology’, *Physical Review Letters* **74**, 846–849.
- Vilenkin, A. (1996*a*), Predictions from Quantum Cosmology, *in* N. Sanchez and A. Zichichi, eds, ‘String gravity and physics at the Planck energy scale. Proceedings of the NATO Advanced Study Institute, Erice, Italy, 8-19 September 1995. Edited by Norma Sanchez and Antonio Zichichi. Dordrecht: Kluwer Academic Publishers, NATO Advanced Study Institutes Series. Volume C476’, p. 345.
- Vilenkin, A. (1996*b*), Quantum Cosmology and the Constants of Nature, *in* K. Sato, T. Sugimotohara and N. Sugiyama, eds, ‘Cosmological Constant and the Evolution of the Universe’, p. 161.
- Wang, L., Caldwell, R. R., Ostriker, J. P. and Steinhardt, P. J. (2000), ‘Cosmic Concordance and Quintessence’, *ApJ* **530**, 17–35.
- Weinberg, S. (1987), ‘Anthropic bound on the cosmological constant’, *Physical Review Letters* **59**, 2607–2610.
- Weinberg, S. (2000), ‘A priori probability distribution of the cosmological constant’, *Phys. Rev. D* **61**(10), 103505.
- Wetherill, G. W. (1996), The Formation Of Habitable Planetary Systems, *in* L. R. Doyle, ed., ‘Circumstellar Habitable Zones’, p. 193.
- Zlatev, I., Wang, L. and Steinhardt, P. J. (1999), ‘Quintessence, Cosmic Coincidence, and the Cosmological Constant’, *Physical Review Letters* **82**, 896–899.