# Simple Waves

## Reference: Landau & Lifshitz, Fluid Mechanics

## **1 Nonlinear waves**

## **1.1 Small amplitude waves**

When we considered sound waves we considered small amplitude perturbations to the fluid equations. In 1D

$$\frac{\partial^2 \delta \rho}{\partial x^2} - \frac{1}{c_s^2} \frac{\partial^2 \delta \rho}{\partial t^2} = 0 \qquad (1)$$

The solution to this equation, of course, is:

$$\delta \rho = f_1(x - c_s t) + f_2(x + c_s t)$$
(2)

where  $f_1$  and  $f_2$  are arbitrary functions. The perturbation to the velocity is given by:

$$\frac{\partial \delta v_x}{\partial t} = -\frac{c_s^2}{\rho} \frac{\partial \delta \rho}{\partial x} = -\frac{c_s^2}{\rho} [f_1'(x - c_s t) + f_2'(x + c_s t)]$$

$$\delta v_x = \frac{c_s}{\rho} [f_1(x - c_s t) - f_2(x + c_s t)]$$
(3)

Note that for the wave travelling in each direction

$$\delta v_x = \pm \frac{c_s}{\rho} \delta \rho \tag{4}$$

Astrophysical Gas Dynamics: Simple Waves

## **1.2 Simple waves**

The above relationship suggests looking for a full nonlinear solution in which

$$v = v(\rho) \tag{5}$$

Such a solution of the Euler equations is called a simple wave. Take the 1D Euler equations

$$\frac{\partial \rho}{\partial t} + \frac{\partial (\rho v)}{\partial x} = 0$$
$$\frac{\partial v}{\partial t} + v \frac{\partial v}{\partial x} + \frac{1}{\rho} \frac{\partial p}{\partial x} = 0$$

Astrophysical Gas Dynamics: Simple Waves

(6)

Since  $v = v(\rho)$  and  $p = p(\rho)$  then p = p(v) and the above equations can be written:

$$\frac{\partial \rho}{\partial t} + \frac{d}{d\rho} (\rho v) \frac{\partial \rho}{\partial x} = 0$$

$$\frac{\partial v}{\partial t} + \left( v + \frac{1}{\rho} \frac{dp}{dv} \right) \frac{dv}{dx} = 0$$

## **1.2.1 Aside - Profile velocity**

We would like to work out the velocity of a point on the density profile, at a fixed density, as shown in the figure. The aim of the

following is to determine  $\frac{\partial x}{\partial x}$ 

$$\left. e \left. \frac{\partial x}{\partial t} \right|_{\rho} \right|_{\rho}$$



To determine the vlocity of a point with constant density, we consider a transformation of independent variables from (x, t) to  $(\rho, t)$  defined by:

$$\rho = \rho(x, t)$$

$$t' = t$$
(8)

## the reverse transformation being:

$$x = x(t', \rho)$$

$$t = t'$$
(9)

In the new variables  $(\rho, t)$ 

$$\frac{\partial \rho}{\partial t'}\Big|_{\rho} = \frac{\partial \rho}{\partial x}\Big|\frac{\partial x}{\partial t'}\Big|_{\rho} + \frac{\partial \rho}{\partial t}\Big|\frac{\partial t}{x\partial t'}\Big|_{\rho}$$
$$= \frac{\partial \rho}{\partial x}\Big|\frac{\partial x}{\partial t'}\Big|_{\rho} + \frac{\partial \rho}{\partial t}\Big|_{x} = 0$$
$$\Rightarrow \frac{\partial x}{\partial t}\Big|_{\rho} = \frac{\frac{\partial \rho}{\partial t}\Big|_{x}}{\frac{\partial \rho}{\partial x}\Big|_{t}}$$

Astrophysical Gas Dynamics: Simple Waves

(10)

Similarly, in order to consider the profle velocity at a given velocity value, we consider a transformation

$$v = v(x, t)$$

$$t' = t$$
(11)

from (x, t) to (v, t) and the rate of change of x at a fixed t is given by

$$\frac{\partial x}{\partial t}\Big|_{v} = -\frac{\frac{\partial v}{\partial t}}{\frac{\partial v}{\partial x}}$$

(12)

#### **1.3 Continued solution for simple wave:**

Given the above expression for a simple wave and the continuity equation we have

$$\frac{\partial x}{\partial t}\Big|_{\rho} = \frac{d}{d\rho}(\rho v) = v + \rho \frac{dv}{d\rho}$$
(13)

The momentum equation implies that

$$\frac{\partial x}{\partial t}\Big|_{v} = v + \frac{1}{\rho}\frac{dp}{dv}$$
(14)

Now the crucial point in the development of this solution is that since  $v = v(\rho)$  then v = constant corresponds to  $\rho = \text{constant}$  and

$$\frac{\partial x}{\partial t}\Big|_{v} = \frac{\partial x}{\partial t}\Big|_{\rho}$$
(15)

Hence,



(16)

#### Astrophysical Gas Dynamics: Simple Waves

Thus *v* as a function of  $\rho$  is given by:

$$v = \pm \int \frac{c_s}{\rho} d\rho \tag{17}$$

We take a polytropic equation of state

$$p = K(s)\rho^{\gamma} \tag{18}$$

and take  $\rho_0$  to be a reference density corresponding to v = 0. Hence,

$$K = p_0 \rho_0^{-\gamma} \qquad c_0^2 = \gamma K \rho_0^{\gamma - 1}$$
$$= \left(\frac{\rho}{\rho_0}\right)^{\gamma} \qquad c_s^2 = c_0^2 \left(\frac{\rho}{\rho_0}\right)^{\gamma - 1} \qquad c_s = c_0 \left(\frac{\rho}{\rho_0}\right)^{\frac{\gamma - \lambda}{2} - \frac{\gamma}{2}}$$

Astrophysical Gas Dynamics: Simple Waves

 $\frac{p}{p_0}$ 

Hence the relationship between V and  $\rho$  for a polytropic gas is:

$$v = \pm \int \frac{c_s}{\rho} d\rho = \pm c_0 \int \left(\frac{\rho}{\rho_0}\right)^{\frac{\gamma - 3}{2}} d\left(\frac{\rho}{\rho_0}\right)$$
(20)  
$$= \pm \frac{2c_0}{\gamma - 1} \left[ \left(\frac{\rho}{\rho_0}\right)^{\frac{\gamma - 1}{2}} - 1 \right] = \pm \frac{2c_0}{\gamma - 1} \left[ \frac{c_s}{c_0} - 1 \right]$$
(20)

Solving for the sound speed:

$$c_s = c_0 \pm \frac{\gamma - 1}{2} v$$
 (21)

Astrophysical Gas Dynamics: Simple Waves

#### **1.3.1 Wave velocity**

$$\left(\frac{\partial x}{\partial t}\right)_{v} = v + \frac{1}{\rho}\frac{dp}{dv} = v + \frac{c_{s}^{2}}{\rho}\frac{d\rho}{dv} = v \pm c_{s}$$
(22)  
Since  $c_{s} = c_{s}(\rho) = c_{s}(v)$  then  
 $x = t(v \pm c_{s}) + f(v)$ (23)

where f(v) is an arbitrary function of v. Using the solution for  $c_s$  then

$$x = t \left[ \frac{\gamma + 1}{2} v \pm c_0 \right] + f(v) \tag{24}$$

This is one of the most useful forms of the solution for a simple wave.

## Astrophysical Gas Dynamics: Simple Waves

NB. Waves travelling to the right represented by the + sign in the above expressions; waves travelling to the left by the - sign.

**1.3.2 Velocity and the formation of shocks** From the above

$$\left(\frac{\partial x}{\partial t}\right)_{v} = \frac{\gamma + 1}{2} v \pm c_{0}$$
(25)

and this shows the nonlinear effect on the profile velocity. When

 $v \ll c_0$  then  $\left(\frac{\partial x}{\partial t}\right)_v = \pm c_0$ , i.e. the standard relation for a sound wave. However, when  $v \sim c_0$  nonlinear effects have a marked effect on the profile.



The increase of the profile velocity with increasing positive v and the decrease with decreasing negative v means that the profile steepens as shown in the figure. The end result is a series of ever steepening profiles which eventually become unphysical.



When the profile becomes infinitely steep a shock forms. In this region the assumptions embodied in the Euler equations break down i.e. it is no longer possible to neglect viscosity. (More about this later.)

At the time of formation of the shock:

$$\left. \frac{\partial x}{\partial v} \right|_{t} = 0 \tag{26}$$

Moreover, the shock represents a point of inflection in the profile so that

$$\left. \frac{\partial^2 x}{\partial v^2} \right|_t = 0 \tag{27}$$

If we use the solution:

$$x = t \left( \pm c_0 + \frac{\gamma + 1}{2} v \right) + f(v)$$
 (28)

then the conditions for the formation of a shock are:

$$\frac{\partial x}{\partial v} = \frac{\gamma + 1}{2}t + f'(v) = 0 \Longrightarrow t = -\frac{2f'(v)}{\gamma + 1}$$
(29)

and

$$f''(v) = 0 (30)$$

Both of these conditions have to be satisfied simultaneously.

**1.3.3 Fluid outside the region at rest** This is a special case.



In this case the condition for a shock to form is simply

$$\frac{\partial x}{\partial v}\Big|_{t} = 0 \quad \text{at} \quad v = 0 \tag{31}$$

since the velocity profile beyond this point is simply v = 0. This gievs the condition  $t = -\frac{2f'(0)}{\gamma + 1}$ 

## 2 Piston moving out of a pipe as an example of a simple wave

## **2.1Formation of solution**



The piston is initially at x = 0 and moves towards the left with acceleration *a*. A sound wave travels towards the right with velocity  $c_0$  since the gas is at rest to the right of the piston.

We can solve for this flow as a simple wave. Take the general solution:

$$x = t \left[ \pm c_0 + \frac{1}{2} (\gamma + 1) v \right] + f(v)$$
(32)

The + sign applies to a wave travelling to the right so that

$$x = t \left[ c_0 + \frac{1}{2} (\gamma + 1) v \right] + f(v)$$
 (33)

(Note: the *wave* is travelling to the right although the piston is travelling to the left.)

The arbitrary function may be calculated using the boundary condition that the gas remains in contact with the piston, i.e.

$$v = -at$$
 at  $x = -\frac{at^2}{2}$  (34)

Write the solution in the form:

$$f(v) = x - t \left[ c_0 + \frac{\gamma + 1}{2} v \right]$$
  

$$\Rightarrow f(-at) = -\frac{at^2}{2} - t \left[ c_0 - \frac{(\gamma + 1)}{2} at \right]$$

$$= \frac{c_0}{a} (-at) + \frac{\gamma}{2a} (-at)^2$$
(35)

#### Astrophysical Gas Dynamics: Simple Waves

Therefore

$$f(v) = \frac{c_0}{a}v + \frac{\gamma}{2a}v^2$$

$$\Rightarrow x = t\left[c_0 + \frac{(\gamma+1)}{2}v\right] + \frac{c_0}{a}v + \frac{\gamma}{2a}v^2$$
(36)

and the solution for *x* as a function of *x* and *t* is given by the solution of the quadratic:

$$\frac{\gamma}{2a}v^2 + \left(\frac{c_0}{a} + \frac{\gamma+1}{2}t\right)v + c_0t - x = 0$$

$$\Rightarrow -v = \frac{c_0}{\gamma} + \frac{\gamma+1}{2\gamma}(at) \pm \sqrt{\left(\frac{c_0}{\gamma} + \frac{\gamma+1}{2\gamma}at\right)^2 + \frac{2a}{\gamma}(x - c_0t)}$$
(37)

Astrophysical Gas Dynamics: Simple Waves

The other boundary condition to incorporate is the condition that a sound wave moves away from the piston. i.e. v = 0 at  $x = c_0 t$ . We can readily see that the - sign in the above expression gives the correct boundary condition so that

$$-v = \frac{c_0}{\gamma} + \frac{\gamma+1}{2\gamma}(at) - \sqrt{\left(\frac{c_0}{\gamma} + \frac{\gamma+1}{2\gamma}at\right)^2 + \frac{2a}{\gamma}(x-c_0t)} \quad (38)$$

Does a shock form in this solution?

$$f(v) = \frac{c_0}{a}v + \frac{\gamma}{2a}v^2$$
$$f'(v) = \frac{c_0}{a} + \frac{\gamma}{a}v$$
$$f''(v) = \frac{\gamma}{a}$$

The second derivative is always non zero so that if a shock forms it does so when v = 0 i.e. when

$$t = -\frac{2f'(0)}{\gamma + 1} = -\frac{2}{\gamma + 1}\frac{c_0}{a} < 0$$
(40)

#### Astrophysical Gas Dynamics: Simple Waves

26/33

(39)

Negative values of *t* are outside the domain of validity of our solution so that a shock does *not* form.

## **2.2Maximum Velocity**

An intriguing aspect of the above solution is that a velocity is reached at which a cavity forms behind the piston. Consider our solution for the sound speed in a simple wave:

$$c_s = c_0 \pm \frac{\gamma - 1}{2} v = c_0 + \frac{\gamma - 1}{2} v \tag{41}$$

Now in order that the density remain positive, the sound speed must also remain positive and

$$c_{0} + \frac{\gamma - 1}{2}v > 0$$

$$-v < \frac{2c_{0}}{\gamma - 1}$$
(42)

Once the gas (and hence the piston) reaches this velocity a vacuum forms.



## **3 Piston moving into gas**



## **3.1 Solution**

Piston accelerates from zero velocity to a velocity u = at and a position  $x = \frac{1}{2}at^2$  at time *t*.

Again we use the general simple wave for a wave moving to the right

$$x = t \left[ c_0 + \frac{\gamma + 1}{2} v \right] + f(v) \tag{43}$$

and we use the boundary condition v = at for gas at the surface of the piston:

$$\Rightarrow \frac{1}{2}at^{2} = t\left[c_{0} + \frac{\gamma + 1}{2}at\right] + f(at)$$
$$\Rightarrow f(at) = -\frac{c_{0}}{a}(at) - \frac{\gamma}{2a}(at)^{2}$$
$$(44)$$
$$f(v) = -\frac{c_{0}}{a}v - \frac{\gamma}{2a}v^{2}$$

Astrophysical Gas Dynamics: Simple Waves

#### **3.2 Formation of shock:**

We have

$$f'(v) = -\frac{c_0}{a} - \frac{\gamma}{a}v$$
 and  $f''(v) = -\frac{\gamma}{a} < 0$  (45)

so that the only place where a shock can form is at v = 0. The time of formation of the shock in this case is:

$$t = -\frac{2f'(0)}{\gamma + 1} = \frac{2}{\gamma + 1} \frac{c_0}{a}$$
(46)

Note that the velocity of the piston is not supersonic with respect to the undisturbed gas when the shock forms:

$$u = at = \frac{2}{\gamma + 1}c_0 < c_0 \text{ for } \gamma > 1$$
 (47)

Astrophysical Gas Dynamics: Simple Waves

## **3.3 Velocity of gas**

The expression for the velocity profile is, in this case (exercise):

$$v = -\left(\frac{c_0}{\gamma} - \frac{\gamma+1}{2\gamma}at\right) + \sqrt{\left(\frac{c_0}{\gamma} - \frac{\gamma+1}{2\gamma}at\right)^2 - \frac{2a}{\gamma}(x - c_0 t)} \quad (48)$$

One can show directly from this expression that the velocity pro-

file becomes infinitely steep at 
$$x = c_0 t$$
 when  $t = \frac{2}{\gamma + 1} \frac{c_0}{a}$ .